9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

45-1089: Rabbit Monoclonal Antibody to His Tag (C-term) (Clone : 25B6E11)

Clonality :	Monoclonal
Clone Name :	25B6E11
Application :	IP,ELISA,ICC/IF,FACS,WB
Format :	Purified
Isotype :	Rabbit IgG
Immunogen Information :	6x His synthetic peptide coupled to KLH

Description

The rabbit immune system generates antibody diversity and optimizes affinity. technology to generate high affinity and specific monoclonal rabbit antibodies. Anti-His Tag (C-term) Antibody (25B6E11), mAb, Rabbit is specific to six histidine tags placed at C-terminal and internal regions of fusion proteins. This antibody can greatly improve the effectiveness of several different kinds of immunoassays, helping researchers identify, detect, and purify polyhistidine fusion proteins in bacteria, insect cells, and mammalian cells. This product is not recommended for N-terminal His-fusion protein detection.

Product Info

Amount :	40 µg
Purification :	Protein A chromatography
Content :	lyophilized with PBS, pH 7.4, containing 0.02% sodium azide. Reconstitute with deionized water or equivalent to 0.5 mg/ml.
Storage condition :	The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.

Application Note

ELISA: 0.05-0.2 µg/ml
Western Blot: 1-2 µg/ml
Immunoprecipitation (IP): 2 µg/ml
Immunocytochemistry/Immunofluorescence (ICC/IF): 1 µg/ml
Flow Cytometry: 1 µg/ml



Figure-1 : Western blot analysis of His Tag (C-term) Antibody (Clone: 25B6E11) at 0.2 Î¼g/ml on 6xHis-tag Fusion Protein, IRDye 800 conjugated Goat anti-rabbit IgG (H&L) was used as secondary Antibody at 0.125 Î¼g/ml.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com



Figure-2 : Western blot analysis of His Tag (C-term) Antibody (Clone: 25B6E11) at 0.2 Î¼g/ml on immunoprecipitation from cell lysates containing His fusion protein, 1: His antibody (4 Î¼g) + His tagged fusion protein, 2: Rabbit IgG (4 Î¼g) + His tagged fusion protein, 3: His tagged fusion protein (Input), IRDye 800 Conjugated Goat anti-rabbit IgG (H&L) was used secondry antibody at 0.125 Î¼g/ml.



Figure-3 : Western blot analysis of cross-reactivity of His Tag (C-term) Antibody (Clone: 25B6E11) at 1Î¼g/ml on cell lysates, A: His Tag Antibody (Clone:25B6E11), B: Rabbit IgG (as Negative control), IRDye 800 Conjugated Goat anti-rabbit IgG (H&L) was used as secondry antibody at 0.125 Î¼g/ml.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.