

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

45-1119: Mouse Monoclonal Antibody to Human C-reactive protein (Clone: 39E5) (Discontinued)

Clonality: Monoclonal Clone Name: 39E5 Application: **ELISA** Purified Format:

Isotype: Mouse IgG1, Kappa Immunogen Information: Human CRP

Description

C-reactive protein (CRP) is an annular, pentameric protein found in blood plasma, in which the levels rise in response to inflammation. It is an acute-phase protein synthesized by the liver that increases following interleukin-6 secretion by macrophages and T cells. CRP binds to the phosphocholine expressed on the surface of dead or dying cells and some bacteria. This activates the complement system, promoting phagocytosis by macrophages, which clears necrotic and apoptotic cells and bacteria.

Product Info

Amount: 40 µg

Purification: Protein A chromatography

Content: 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody Storage condition:

can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below.

Avoid repeated freezing and thawing cycles.

Application Note

ELISA detection: 0.02-0.2 µg/ml

Western blot: 1-2 µg/ml




Figure-1: ELISA binding of CRP Antibody (Clone: 39E5) with Human CRP, Coating antigen: Human CRP at 1 µg/ml, Human CRP antibody dilution start from 2000 ng/ml, EC50=11.58 ng/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2 : Western blot analysis of CRP Antibody (Clone: 39E5) at 1 μ g/ml on Human CRP (1-5: 100 ng, 50 ng, 10 ng, 5 ng & 1 ng respectively), IRDye 800 conjugated Goat anti-Mouse IgG was used as Secondary Antibody at 0.125 μ g/ml.