

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-1089: Anti-Carcinoembryonic Antigen (CEA) / CD66 Recombinant Mouse Monoclonal Antibody (Clone:C66/1983R)

Monoclonal Clonality: Clone Name: C66/1983R

Application: IHC Reactivity: Human

CEACAM1 & CEACAM5 Gene:

1048 & 634 Gene ID: **Uniprot ID:** P06731 Format: Purified

Carcinoembryonic Antigen-related Cell Adhesion Molecule 5, CEACAM5, CD66, Biliary **Alternative Name:**

Glycoprotein (BGP-1)

Isotype: Rabbit IgG

Immunogen Information: Human colon carcinoma extract

Description

This antibody recognizes proteins of 80-200kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MAb does not react with nonspecific cross-reacting antigen (NCA) and with human polymorphonuclear leucocytes. It shows no reaction with a variety of normal tissues and is suitable for staining of formalin/paraffin tissues. CEA is not found in benign glands, stroma, or malignant prostatic cells. Antibody to CEA is useful in detecting early foci of gastric carcinoma and in distinguishing pulmonary adenocarcinomas (60-70% are CEA+) from pleural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is seen in adenocarcinomas from the lung, colon, stomach, esophagus, pancreas, gallbadder, urachus, salivary gland, ovary, and endocervix.

Product Info

Amount: 20 μg / 100 μg Purification: Protein A/G

200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Content:

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is Storage condition:

stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded Human Melanoma stained with CEA Rabbit Recombinant Monoclonal Antibody (C66/1983R).




Figure 2: SDS-PAGE Analysis of Purified CEA Rabbit Recombinant Monoclonal Antibody (C66/1983R).