

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-1213: Anti-Moesin Recombinant Mouse Monoclonal Antibody (Clone:rMSN/492)

Clonality: Monoclonal **Clone Name:** rMSN/492 Application: FACS.IF.WB.IHC Reactivity: Human, Rat

Gene: MSN Gene ID: 4478 **Uniprot ID:** P26038 Format: **Purified**

Membrane-organizing extension spike protein; Moesin/anaplastic lymphoma kinase fusion **Alternative Name:**

protein; MSN/ALK fusion

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant full-length human Moesin protein

Description

Recognizes 78kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the submembranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibroblastic, endothelial, epithelial, and neuronal cell lines but not in blood cells. The ERM proteins, ezrin, radixin, and moesin are involved in a variety of cellular functions, such as cell adhesion, migration, and the organization of cell surface structures, and are highly homologous, both in protein sequence and in functional activity, with merlin/schwannomin, a neurofibromatosis-2-associated tumor-suppressor protein. Cell lines of epithelial and mesothelial origin contain both moesin and radixin whereas cells of endothelial and lymphoid origin express moesin.

Product Info

Amount: 20 μg / 100 μg **Purification:** Protein A/G

200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Content:

0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2Ă∏Âμg/million cells); Immunofluorescence (2-4Ã∏Âμg/ml); Western Blot (1-2Ã∏Âμg/ml); Immunohistochemistry (Formalin-fixed) (1-2̸µq/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded human Placenta stained with Moesin Mouse Recombinant Monoclonal Antibody (rMSN/492).




Figure 2: SDS-PAGE Analysis Purified Moesin Mouse Recombinant Monoclonal Antibody (rMSN/492).