

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-1235: Anti-Parathyroid Hormone (PTH) (N-Terminal) Recombinant Rabbit Monoclonal Antibody (Clone:PTH/2295R)

Clonality: Monoclonal Clone Name: PTH/2295R

Application: IHC
Reactivity: Human
Gene: PTH
Gene ID: 5741
Uniprot ID: P01270
Format: Purified

Alternative Name: hPTH; Parathormone; Parathyrin; Parathyroid hormone 1 (PTH1); Parathyroid hormone (PTH)

Isotype: Rabbit IgG

Immunogen Information: Synthetic peptide corresponding to amino acids around 1 to 34 of mature PTH conjµgated to

a carrier

Description

Epitope of this MAb maps in between aa 1-34. PTH is a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood Ca2+ levels by dissolving the salts in bone and preventing their renal excretion. It is produced in the parathyroid gland as an 84 amino acid single chain polypeptide. It can also be secreted as N-terminal truncated fragments or C-terminal fragments after intracellular degradation, as in case of hypercalcemia. Defects in this gene are a cause of familial isolated hypoparathyroidism (FIH); also called autosomal dominant hypoparathyroidism or autosomal dominant hypocalcemia. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures, tetany and cramps. FIH exist both as autosomal dominant and recessive forms of hypoparathyroidism.

Product Info

Amount : $20 \mu g / 100 \mu g$ Purification : Protein A/G

Content: 200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2 \tilde{A} \square \hat{A} μ g/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded human Parathyroid stained with PTH Rabbit Recombinant Monoclonal Antibody (PTH/2295R).




Figure 2: SDS-PAGE Analysis Purified PTH Rabbit Recombinant Monoclonal Antibody (PTH/2295R).