

## 12-1081: Anti-Bcl-6 (Follicular Lymphoma Marker) Recombinant Rabbit Monoclonal Antibody (Clone:BCL6/1951R)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal   |
| <b>Clone Name :</b>            | BCL6/1951R   |
| <b>Reactivity :</b>            | Human  |
| <b>Gene :</b>                  | BCL6   |
| <b>Gene ID :</b>               | 604  |
| <b>Uniprot ID :</b>            | P41182   |
| <b>Format :</b>                | Purified   |
| <b>Alternative Name :</b>      | B-cell lymphoma 5 protein; B-Cell Lymphoma 6 Protein; BCL5; BCL6; BCL6A; cys his2 zinc finger transcription factor; Lymphoma Associated Zinc Finger Gene On Chromosome 3 (LAZ3); Zinc finger and BTB domain-containing protein 27 (ZBTB27); Zinc Finger Protein 51 (ZNF51); zinc finger transcription factor BCL6S |
| <b>Immunogen Information :</b> | Recombinant fragment of human bcl-6 protein (around aa 256-389) (exact sequence is proprietary)  |

### Description

Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large (L&H) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 20 µg / 100 µg  |
| <b>Purification :</b>      | Protein A/G   |
| <b>Content :</b>           | 200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| <b>Storage condition :</b> | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.               |




Figure 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Bcl-6 Rabbit Recombinant Monoclonal Antibody (BCL6/1951R).




Figure 2: SDS-PAGE Analysis of Purified Bcl-6 Rabbit Recombinant Monoclonal Antibody (BCL6/1951R).