

12-1247: Anti-Thrombomodulin / CD141 (Endothelial Cell Marker) Recombinant Mouse Monoclonal Antibody (Clone:rTHBD/1591)

Clonality :	Monoclonal
Clone Name :	rTHBD/1591
Application :	ELISA,IHC
Reactivity :	Human
Gene :	THBD
Gene ID :	7056
Uniprot ID :	P07204
Format :	Purified
Alternative Name :	AHUS6; BDCA3; CD141; Fetomodulin; Thbd; THPH12; THRM; Thrombomodulin (TM)
lsotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant fragment of human Thrombomodulin protein (around aa 69-194) (exact sequence is proprietary)

Description

It recognizes a protein of 75kDa, identified as Thrombomodulin. Thrombomodulin is a transmembrane glycoprotein with natural anticoagulant properties. It is normally expressed by a restricted number of cells, such as endothelial and mesothelial cells. In addition, synovial lining and syncytio-trophoblasts of placenta also express thrombomodulin. This protein is present in almost all of benign vascular tumors and majority of malignant vascular tumors (Kaposi s sarcoma, angiosarcoma, and epithelioid hemangioendothelioma). Hence, anti-thrombomodulin serves as a sensitive marker for lymphatic endothelial cells and their tumors. Recently, thrombomodulin antibody has been used for mesothelial cells and malignant mesotheliomas.Â

Product Info

Amount :	20 μg / 100 μg
Purification :	Protein A/G
Content :	$200\mu g/ml$ of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (Use Ab at 2-4Ã^[]µg/ml for coating) (Order Ab without BSA); Immunohistochemistry (Formalin-fixed) (1-2Ã^[]µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)

∗ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 1: Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Thrombomodulin / CD141 Monoclonal Antibody (rTHBD/1591).



Figure 2: Formalin-fixed, paraffin-embedded human Placenta stained with Thrombomodulin / CD141 Monoclonal Antibody (rTHBD/1591).

Figure 3:SDS-PAGE Analysis Purified Thrombomodulin Mouse Recombinant Monoclonal Antibody (rTHBD/1591).



Figure 4:Analysis of Protein Array containing >19,000 full-length human proteins using Thrombomodulin Recombinant Mouse Monoclonal Antibody (rTHBD/1591) Zand S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Zscores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.