# **w** abeomics

### 12-1261: Anti-Tyrosinase (Melanoma Marker) Recombinant Mouse Monoclonal Antibody (Clone:rOCA1/812)

Clonality :	Monoclonal
Clone Name :	rOCA1/812
Application :	IHC
Reactivity :	Human
Gene :	TYR
Gene ID :	7299
Uniprot ID :	P14679
Format :	Purified
Alternative Name :	ATN, CMM8, LB24-AB, Monophenol monooxygenase, OCA1, OCA1A, Oculocutaneous albinism IA, SHEP3, SK29-AB, Tumor rejection antigen AB, TYR
Isotype :	Mouse IgG1, kappa
Immunogen Information : Recombinant full-length human tyrosinase protein	

#### Description

Recognizes a cluster of proteins between 70-80kDa, identified as tyrosinase. Occasionally a minor band at 55kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-containing metalloglycoprotein that catalyzes several steps in the melanin pigment biosynthetic pathway; the hydroxylation of tyrosine to L-3,4-dihydroxy-phenylalanine (dopa), and the subsequent oxidation of dopa to dopaquinone. Mutations of the tyrosinase gene occur in various forms of albinism. Tyrosinase is one of the targets for cytotoxic T-cell recognition in melanoma patients. Staining of melanomas with this MAb shows tyrosinase in melanotic as well as amelanotic variants.This MAb is a useful marker for melanocytes and melanomas.

#### **Product Info**

Amount :	20 µg / 100 µg
Purification :	Protein A/G
Content :	$200\mu$ g/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### **Application Note**

Immunohistochemistry (Formalin-fixed) (1-2Ã µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 &degC followed by cooling at RT for 20 minutes),

## **₩** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded Melanoma stained with Tyrosinase Mouse Recombinant Monoclonal Antibody (rOCA1/812).

Figure 2: SDS-PAGE Analysis Purified Tyrosinase Mouse Recombinant Monoclonal Antibody (rOCA1/812).