

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-1085: Anti-Caldesmon, HMW (h-Caldesmon) (Smooth Muscle Marker) Recombinant Mouse Monoclonal Antibody (Clone:rCALD1/820)

Clonality: Monoclonal **Clone Name:** rCALD1/820 Application: IHC,WB Reactivity: Human Gene: CALD1 Gene ID: 800 **Uniprot ID:** 005682 Format: **Purified** 

Alternative Name: CAD; CALD1; Caldesmon 1 Isoform 1; Caldesmon 1 Isoform 2; Caldesmon 1 Isoform 3;

Caldesmon 1 Isoform 4; Caldesmon 1 Isoform 5; CDM; HCAD; LCAD; NAG22

**Isotype:** Mouse IgG1, kappa

Immunogen Information: Recombinant full-length human CALD1 protein

## **Description**

Recognizes a protein of 150kDa, which is identified as the high molecular weight variant of Caldesmon. Two closely related variants of human caldesmon have been identified which are different in their electrophoretic mobility and cellular distribution. The h-caldesmon variant (120-150kDa) is predominantly expressed in smooth muscle whereas l-caldesmon (70-80kDa) is found in non- muscle tissue and cells. Neither of the two variants has been detected in skeletal muscle. This MAb recognizes only the 150kDa variant (h-caldesmon) in Western blots of human aortic media extracts and is unreactive with fibroblast extracts from cultivated human foreskin. Caldesmon is a developmentally regulated protein involved in smooth muscle and non-muscle contraction.

## **Product Info**

Amount :  $20 \mu g / 100 \mu g$ Purification : Protein A/G

Content: 200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Western Blot (1-2 $\mu$ g/ml); ,Immunohistology (Formalin-fixed) (0.25-0.5 $\mu$ g/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 1mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes),



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded human Leiomyosarcoma stained with Caldesmon Mouse Recombinant Monoclonal Antibody (rCALD1/820).




Figure 2: SDS-PAGE Analysis of Purified Caldesmon Mouse Recombinant Monoclonal Antibody (rCALD1/820).