9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

12-1171: Anti-E-Cadherin (CDH1) / CD324 (Intercellular Junction Marker) Recombinant Rabbit Monoclonal Antibody (Clone:CDH1/2208R)

Clonality :	Monoclonal
Clone Name :	CDH1/2208R
Application :	IHC,FACS,WB,IF
Reactivity :	Human
Gene :	CDH1
Gene ID :	999
Uniprot ID :	P12830
Format :	Purified
Alternative Name :	Arc 1; cadherin 1 type 1 E-cadherin; Cadherin1; CAM 120/80; CD324; CDH1; CDHE; E-Cad/CTF3; E-cadherin; ECAD; Epithelial cadherin; epithelial calcium dependent adhesion protein; Liver cell adhesion molecule (LCAM); Uvomorulin (UVO)
Isotype :	Rabbit IgG
Immunogen Information :	Recombinant full-length human E-Cadherin protein

Description

Recognizes a protein of 120-80kDa, identified as E-cadherin. Cadherins comprise a family of Ca2+-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- andP-cadherin, consist of large extracellular domains characterized by a series of five homologous NH2 terminal repeats. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as catenin, to regulate cadherin function.E-cadherin plays an important role in epithelial cell adhesion. A decreased expression of E-cadherin is associated with metastatic potential and poor prognosis in breast cancer, prostate and esophageal cancer. In combination with p120 Catenin, it is useful for the differentiation between ductal (E-cadherin +) and lobular (E-cadherin -) breast carcinomas. It may also help in diagnosis of mesothelioma.

Product Info

Amount :	20 µg / 100 µg
Purification :	Protein A/G
Content :	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2µg/million cells);Immunofluorescence (1-2µg/ml); Western Blot(1-2µg/ml); ,Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded human Cervix stained with E-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH1/2208R).




Figure 2: SDS-PAGE Analysis of Purified E-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH1/2208R).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.