

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-1164: Anti-Cytokeratin, Basic (Type II or HMW) Recombinant Mouse Monoclonal Antibody (Clone:rKRTH/2148)

Clonality: Monoclonal Clone Name: rKRTH/2148

Application: IHC

Reactivity: Human, Mouse, Rat

 Gene :
 KRT76

 Gene ID :
 51350

 Uniprot ID :
 Q01546

 Format :
 Purified

Alternative Name: KRT2B; KRT2P; HUMCYT2A; Keratin, type II Cytoskeletal 2 oral; K76; Keratin 2p (K2P);

Keratin-76; Cytokeratin-2P (CK-2P; Type-II Keratin Kb9

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant full-length of human KRT76 protein

Description

This antibody recognizes CK1, CK5, CK10 and CK14. In normal epithelia, it stains stratified epithelia, myoepithelial cells and basal cells in the prostate gland and bronchi. This MAb shows no reactivity with hepatocytes, pancreatic acinar cells, proximal renal tubules, or endometrial glands; there is no reactivity with cells derived from simple epithelia. Mesenchymal tumors, lymphomas, melanomas, neural tumors, and neuroendocrine tumors are negative with this antibody. It stains myoepithelial cells and has been shown to be useful in distinguishing prostate adenocarcinoma from benign prostate. This antibody has also been useful in separating benign from malignant intraductal breast proliferations.

Product Info

Amount : 20 μg / 100 μg **Purification :** Protein A/G

Content: 200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2 \tilde{A} \square $\hat{A}\mu$ g/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 \tilde{A} \square \hat{A} °C followed by cooling at RT for 20 minutes).




Figure 1: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with CK HMW Mouse Recombinant Monoclonal Antibody (rKRTH/2148).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: SDS-PAGE Analysis of Purified CK HMW Mouse Recombinant Monoclonal Antibody (rKRTH/2148).