

#### 44-1058: Anti-p16 Monoclonal Antibody (Clone:IHC016)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	IHC016
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	CDKN2A
<b>Gene ID :</b>	1029
<b>Uniprot ID :</b>	P42771
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CDKN2A,CDKN2,MTS1
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Recombinant human p16INK4A C-terminus protein

#### Description

The p16 (p16INK4A) protein is a cyclin-dependent kinase inhibitor that plays an important regulatory role in the cell cycle. By controlling the transition between the G1 and S phases through regulation of retinoblastoma protein, p16 decelerates cellular differentiation and therefore acts as a tumor suppressor, making it the key marker in several human cancers including head and neck cancer, perianal lesions, melanomas, gliomas, lymphomas, and some types of leukemia. p16 is also clinically indicated in carcinomas of the esophagus, pancreas, lung, biliary tract, liver, colon, and urinary bladder.

#### Product Info

<b>Amount :</b>	0.1 ml / 0.5 ml
<b>Purification :</b>	Protein A/G Chromatography
<b>Content :</b>	Tris Buffer, pH 7.3 - 7.7, with 1% BSA and <0.1% Sodium Azide
<b>Storage condition :</b>	Store at 2°C - 8°C. Do not freeze.

#### Application Note

Recommended dilutions: Immunohistochemical analysis: 1:100 - 1:200. However, this need to be optimized based on the research applications.