

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1545: Monoclonal Antibody to CD31 / PECAM-1 (Endothelial Cell Marker)(C31.3 + JC/70A)(Discontinued)

Clone Name: Monoclonal
Clone Name: C31.3 + JC/70A
Application: FACS,IF,IHC

Reactivity: Human,Cynomolgus Monkey,Rabbit

Gene : PECAM1
Gene ID : 5175
Uniprot ID : P16284
Format : Purified
Alternative Name : PECAM1
Isotype : Mouse IgG

Immunogen Information: Human recombinant CD31 protein (C31.3) & Membrane preparation of a spleen from a

patient with hairy cell leukemia (JC/70A)

Description

CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 MAb reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.

Product Info

Amount : 100 μg

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly

toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.


Application Note

Flow Cytometry (0.5-1 \tilde{A} | \hat{A} µg/million cells in 0.1ml); Immunofluorescence (0.5-1 \tilde{A} | \hat{A} µg/ml); Immunohistology (Formalin-fixed) (1-2 \tilde{A} | \hat{A} µg/ml for 30 minutes at RT); (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes); Optimal dilution for a specific application should be determined.




9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CD31 Monoclonal Antibody (C31.3 \pm JC/70A)



Formalin-fixed, paraffin-embedded human Angiosarcoma stained with CD31 Monoclonal Antibody (C31.3 \pm JC/70A)