

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1685: APC Conjugated, Anti-CD11a / LFA-1 alpha chain Monoclonal Antibody (Clone:MEM-25)

Clonality: Monoclonal Clone Name: MEM-25 Application: **FACS** Reactivity: Human Conjugate: APC **ITGAL** Gene: Gene ID: 3683 **Uniprot ID:** P20701

Alternative Name: ITGAL,CD11A Isotype: Mouse IgG1

Immunogen Information: Leukocytes from a pacient suffering from a LGL-type leukaemia.

Description

CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity conformation. Cell stimulation by chemokines or other signals leads to induction the high-affinity conformation, which supports tight binding of LFA-1 to its ligands, the intercellular adhesion molecules ICAM-1, -2, -3. LFA-1 is thus involved in interaction of various immune cells and in their tissue-specific settlement, but participates also in control of cell differentiation and proliferation and of T-cell effector functions. Blocking of LFA-1 function by specific antibodies or small molecules has become an important therapeutic approach in treatment of multiple inflammatory diseases. For example, humanized anti-LFA-1 antibody Efalizumab (Raptiva) is being used to interfere with T cell migration to sites of inflammation; binding of cholesterol-lowering drug simvastatin to CD11a allosteric site leads to immunomodulation and increase in lymphocytic cholinergic activity.

Product Info

Amount: 100 tests

Storage condition : Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of human peripheral blood cells with anti-human CD11a (MEM-25) APC.