

### 30-1086: Anti-CD11b activation epitope Monoclonal Antibody (Clone:CBRM1/5)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CBRM1/5
<b>Application :</b>	IP, FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	Unconjugated
<b>Gene ID :</b>	3684
<b>Uniprot ID :</b>	P11215
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Mac-1, Integrin alpha M, ITGAM, CR3A, MO1A, MAC1A
<b>Isotype :</b>	Mouse IgG1 kappa
<b>Immunogen Information :</b>	Information not available

#### Description

CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.<br>

**Specificity:** The mouse monoclonal antibody CBRM1/5 recognizes an activation-dependent epitope on extracellular part of CD11b (Mac-1alpha), a 165-170 kDa type 1 transmembrane protein mainly expressed on monocytes, granulocytes and NK-cells. The antibody recognizes a subset of CD11b molecules on neutrophils and monocytes activated with chemoattractants or phorbol esters and it does not recognize CD11b on non-activated cells.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography.
<b>Content :</b>	Concentration: 1 mg/ml Formulation: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

#### Application Note

Flow cytometry: Recommended dilution: 1-4 µg/ml

Flow cytometry: Recommended dilution: 1-4 µg/ml