

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1005: Anti-CD16 / FcgammaRIII Monoclonal Antibody (Clone:MEM-154)-Azide free

Clonality: Monoclonal
Clone Name: MEM-154
Application: FACS
Reactivity: Human
Gene: FCGR3A
Gene ID: 2214
Uniprot ID: P08637

Alternative Name: FCGR3A,CD16A,FCG3,FCGR3,IGFR3

Isotype: Mouse IgG1

Immunogen Information: Human granulocytes

Description

CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution: 5-10 μg/ml Positive control: PBL (peripheral blood lymphocytes)

Application note: The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. **Immunoprecipitation Western Blotting** Application note: Non-reducing conditions. **Functional Application** The antibody MEM-154 blocks binding of human IgG to FcgammaRIII.




Figure 1: Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD16 (MEM-154) purified antibody (azide free, concentration in sample 2 $\hat{1}\frac{1}{4}$ g/ml) GAM APC.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of human neutrophil granulocytes (red-filled) from CD16 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD16 (MEM-154) purified antibody (azide free, concentration in sample 2 $\hat{l}^{1}/4$ g/ml) GAM APC.