

## 30-1120: Anti-CD16 / FcgammaRIII Monoclonal Antibody (Clone:MEM-154)

| Clonality :                                | Monoclonal                    |
|--|-------------------------------|
| Clone Name :                               | MEM-154                       |
| Application :                              | FACS                          |
| Reactivity :                               | Human                         |
| Gene :                                     | FCGR3A                        |
| Gene ID :                                  | 2214                          |
| Uniprot ID :                               | P08637                        |
| Format :                                   | Purified                      |
| Alternative Name :                         | FCGR3A,CD16A,FCG3,FCGR3,IGFR3 |
| Isotype :                                  | Mouse IgG1                    |
| Immunogen Information : Human granulocytes |                               |

#### Description

CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

#### **Product Info**

| Amount :            | 0.1 mg  |
|---------------------|---|
| Purification :      | Purified by protein-A affinity chromatography |
| Storage condition : | Store at 2-8°C. Do not freeze.                |

### **Application Note**

Flow Cytometry Recommended dilution: 5-10 µg/ml

Positive control: PBL (peripheral blood lymphocytes)

Application note: The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. Immunoprecipitation Western Blotting Application note: Non-reducing conditions. Functional Application The antibody MEM-154 blocks binding of human IgG to FcgammaRIII.

# **∗** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 1: Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD16 (MEM-154) purified antibody (concentration in sample 2  $\hat{1}$ /4g/ml) GAM APC.

Figure 2: Separation of FasL transfected L5178Y cells stained using anti-human CD178 (NOK-1) purified antibody (concentration in sample 9  $\hat{I}_{4}$ g/ml, GAM APC, red-filled) from FasL transfected L5178Y cells unstained by primary antibody (GAM APC, black-dashed) in flow cytometry analysis (surface staining).