

## 30-2066: Anti-CD16 / FcgammaRIII Monoclonal Antibody (Clone:3G8)-FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	3G8
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Gene :</b>	FCGR3A
<b>Gene ID :</b>	2214
<b>Uniprot ID :</b>	P08637
<b>Alternative Name :</b>	FCGR3A,CD16A,FCG3,FCGR3,IGFR3
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Human neutrophils

### Description

CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

### Product Info

<b>Amount :</b>	100 tests
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD16 (3G8) FITC antibody (4  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).




Figure 2: Flow cytometry multicolor surface staining pattern of human peripheral whole blood stained using anti-human CD16 (3G8) FITC antibody (4  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood) and anti-human CD3 (UCHT1) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).




Figure 3: Separation of human CD16 positive CD3 negative lymphocytes (red-filled) from CD16 negative CD3 positive lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD16 (3G8) FITC antibody (4  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).