

## 30-2224: Anti-CD16 / Fcγ<sub>3</sub> Monoclonal Antibody (Clone:LNK16)-PE Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	LNK16
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Gene :</b>	FCGR3A
<b>Gene ID :</b>	2214
<b>Uniprot ID :</b>	P08637
<b>Alternative Name :</b>	FCGR3A,CD16A,FCG3,FCGR3,IGFR3
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Normal human peripheral blood granulocytes

### Description

CD16 (Fcγ<sub>3</sub>) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human Fcγ<sub>3</sub> is expressed in two forms - Fcγ<sub>3</sub>-A and -B. Fcγ<sub>3</sub>-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεRI-γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell Fcγ<sub>3</sub>-A is associated, moreover, with FcεRI-β subunit. Besides IgG, Fcγ<sub>3</sub>-A can be triggered also by oligomeric IgE. Fcγ<sub>3</sub>-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

### Product Info

<b>Amount :</b>	100 tests
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD16 (LNK16) PE antibody (concentration in sample 9 µg/ml).




Figure 2: Expression profiling on peripheral blood subsets using Anti-human CD16 PE antibody (clone LNK16). Adaptive panel




Figure 3: Expression profiling on peripheral blood subsets using Anti-human CD16 PE antibody (clone LNK16). Innate panel




Figure 4: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD3 (UCHT1) APC antibody (10 µl reagent / 100 µl of peripheral whole blood) and anti-human CD16 (LNK16) PE antibody (concentration in sample 9 µg/ml).




Figure 5: Separation of human neutrophil granulocytes (red-filled) from CD3 positive CD16 negative T cells (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD16 (LNK16) PE antibody (concentration in sample 9 µg/ml).