

30-1789: Anti-CD19 Monoclonal Antibody (Clone:4G7)-APC Conjugated

| Clonality : | Monoclonal |
|--|-------------------------|
| Clone Name : | 4G7 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | APC |
| Gene : | CD19 |
| Gene ID : | 930 |
| Uniprot ID : | P15391 |
| Alternative Name : | CD19, B4, Leu-12, CVID3 |
| Isotype : | Mouse IgG1 |
| Immunogen Information : Human CCL (chronic lymphocytic leukemia) cells | |

Description

Specificity: The mouse monoclonal antibody 4G7 recognizes an extracellular epitope of human CD19.

CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell treshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

Product Info

| Amount : | 100 tests |
|---------------------|---|
| Purification : | Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Content : | Formulation : Stabilizing phosphate-buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage condition : | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure 1: Surface staining of human peripheral blood leukocytes with anti-human CD19 (4G7) APC.