

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1081: Anti-CD267 / TACI Monoclonal Antibody (Clone:1A1)

Clonality: Monoclonal

Clone Name: 1A1
Application: FACS
Reactivity: Human
Gene: TNFRSF13B
Gene ID: 23495
Uniprot ID: 014836
Format: Purified

Alternative Name: TNFRSF13B,TACI Isotype: Rat IgG2a

Immunogen Information: CD267-transfected RBL cells

Description

CD267 / TACI (Transmembrane Activator Calcium modulator and cyclophilin ligand Interactor), a TNFR superfamily transmembrane protein, is expressed on B cells (predominantly on CD27+ memory cells), multiple myeloma cells and B cell chronic lymphocytic leukemia (B-CLL). Its triggering leads to activation of the transcription factors NFAT, AP1, and NF-kappa-B. It plays a crucial role in humoral immunity. Mutations in CD267 are associated with common variable immunodeficiency and IgA deficiency.

Product Info

Amount: 0.1 mg

Purification: Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD267 (1A1) purified antibody (concentration in sample 3 µg/ml) GAM APC.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com





Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD19 (LT19) PE antibody (20 μ l reagent / 100 μ l of peripheral whole blood) and anti-human CD267 (1A1) purified antibody (concentration in sample 3 μ g/ml) GAM APC.



CD267 purified / GAM APC

Figure 3: Separation of human CD267 positive CD19 positive B cells (red-filled) from CD267 negative CD19 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using antihuman CD267 (1A1) purified antibody (concentration in sample 3 μ g/ml) GAM APC.