

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1406: Anti-CD326 / EpCAM Monoclonal Antibody (Clone:323/A3)

Clonality: Monoclonal **Clone Name:** 323/A3 FACS Application: Reactivity: Human Gene: **EPCAM** Gene ID: 4072 **Uniprot ID:** P16422 **Purified** Format:

Alternative Name : EPCAM,GA733-2,M1S2,M4S1,MIC18,TACSTD1,TROP1

Isotype: Mouse IgG1

Immunogen Information: Human breast cancer MCF-7 cells

Description

CD326 / EpCAM (also known as ESA, EGP40, EGP-2, KSA1/4, CO17-1A, GA733-2, MOC31, Ber-EP4) is a 40 kDa transmembrane glycoprotein serving as adhesion molecule in the basolateral membranes in a variety of epithelial cells. CD326 mediates calcium-independent homotypic cell-cell adhesions. CD326 over-expression has been detected in many epithelial tumours and is often associated with bad prognosis. It has been used for diagnostics of (pre-) malignancies at early stages.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Western blotting: Non-reducing conditions; recommended dilution: 1-2 µg/ml; positive control: MCF-7 cells.

Immunoprecipitation: Recommended dilution: 1-4 μg / ml per 100-500 μg of protein in 1 ml lysate.

Immunohistochemistry (paraffin sections): Recommended dilution: 1-10 μ g/ml; pretreatment: tissue section digestion with pepsin (15 min at RT or 10 min at 37°C, 1 mg / ml Tris-HCl, pH 2.0); positive tissue: breast carcinoma.

pepsin (15 min at R1 or 10 min at 37°C, 1 mg / mi 11is-HCl, pH 2.0); positive tissue: breast ca Immunocytochemistry: Recommended dilution: 1-10 μg/ml.

Flow cytometry: Recommended dilution: 1-4 µg/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-1: Western blotting analysis of human CD326 (EpCAM) using mouse monoclonal antibody 323/A3 on lysates of MCF-7 cell line and HEK293T/17 cell line (CD326 non-expressing cell line; negative control) under non-reducing and reducing conditions. Nitrocellulose membrane was probed with 2 $\mu g/ml$ of mouse anti-CD326 monoclonal antibody 323/A3 followed by IRDye800-conjugated anti-mouse IgG1 secondary antibody. A specific band was detected for CD326 protein at approximately 40 kDa.