

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2471: Anti-CD38 Monoclonal Antibody (Clone:HIT2)-PE/DyLight 594 Conjugated (Discontinued)

Clonality: Monoclonal

Clone Name: HIT2 Application: **FACS** Reactivity: Human Gene: CD38 Gene ID: 952 **Uniprot ID:** P28907 **Alternative Name:** CD38 Mouse IgG1 Isotype:

Immunogen Information: Human thymocytes in foetus

Description

CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by enzymatic activity of its extracellular part. CD38 acts as NAD+ glycohydrolase converting NAD+ into ADP-ribose, as ADP-ribosyl cyclase producing cADPR and as cADPR hydrolase, thus affecting levels of calcium-mobilizing metabolites. ADPR produced by CD38 serves as an important second messenger of neutrophil and dendritic cell migration.

Product Info

Amount: 100 tests

Purified antibody is conjugated with activated tandem dye of R-phycoerythrin-DyLight®594 (PE-

Purification: DyLight®594) under optimum conditions and unconjugated antibody and free fluorochrome are

removed by size-exclusion chromatography.

Content: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide **Storage condition:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD38 (HIT2) PE-DyLight® 594 (4 μl reagent / 100 μl of peripheral whole blood).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of human CD38 positive monocytes (red-filled) from human CD38 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD38 (HIT2) PEDyLight \$ 594 (4 μ l reagent / 100 μ l of peripheral whole blood).