

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1800: Anti-CD4 Monoclonal Antibody (Clone:OX-35)-APC Conjugated

Clonality: Monoclonal Clone Name: OX-35

Application: FACS, IHC, ICC

Reactivity: Rat

Conjugate: APC

Gene: Cd4

Gene ID: 24932

Uniprot ID: P05540

Alternative Name: Cd4

Isotype: Mouse IgG2a

Immunogen Information: MLR generated rat Th cells

Description

CD4 is a single chain transmembrane glycoprotein of immunoglobulin supergene family. In its extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). The intracellular region of CD4 associates with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. CD4 binds to MHC class II molecules (by CDR2-like region in CD4 domain 1), HIV envelope protein gp120 (by CDR2-like region in CD4 domain 1) and other ligands, such as IL-16 (by to CD4 domain 3) or L-selectin. CD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection. CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differenciation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

Product Info

Amount: 0.1 mg

Storage condition : Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.