

30-2362: Anti-CD42a Monoclonal Antibody (Clone:GR-P)-PE Conjugated

Clonality :	Monoclonal
Clone Name :	GR-P
Application :	FACS
Reactivity :	Human
Conjugate :	PE
Gene :	GP9
Gene ID :	2815
Uniprot ID :	P14770
Alternative Name :	GP9
Isotype :	Mouse IgG1
Immunogen Information :	Human acute lymphoblastic leukemia cells

Description

Specificity: The mouse monoclonal antibody GR-P (also known as GRP-P) recognizes an extracellular epitope of CD42a (glycoprotein 9), a 22 kDa transmembrane protein constitutively expressed on megakaryocytes and platelets.

Description: CD42a, also known as Glycoprotein 9 (GPIX), composes together with GPIb alpha, GPIb beta and GPV the GPIb-IX-V receptor complex critical in the process of platelet-rich thrombus formation by tethering the platelet to a thrombogenic surface. CD42b binds to von Willebrand factor (VWF) exposed at a site of vascular injury, as well as to thrombin, coagulation factors XI and XII, high molecular weight kininogen, TSP-1, integrin Mac-1 and P-selectin. Defects in the gene encoding CD42a are a cause of Bernard-Soulier syndrome, also known as giant platelet disease. These patients have unusually large platelets and have a clinical bleeding tendency.

Product Info

Amount :	100 tests
Purification :	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD42a (GR-P) PE antibody (10 µl reagent / 100 µl of peripheral whole blood).




Figure 2: Separation of human thrombocytes (red-filled) from human neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD42a (GR-P) PE antibody (10 µl reagent / 100 µl of peripheral whole blood).