

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1096: Anti-CD7 Monoclonal Antibody (Clone:124-1D1)

Clonality: Monoclonal **Clone Name:** 124-1D1 Application: FACS Reactivity: Human Gene: CD7 Gene ID: 924 **Uniprot ID:** P09564 Format: **Purified Alternative Name:** CD7

Isotype: Mouse IgG1 **Immunogen Information:** not available

Description

CD7, also known as gp40, is a member of the immunoglobulin superfamily found on T cells, NK cells, thymocytes, hematopoietic progenitors, and monocytes (weakly). CD7 is also expressed on acute lymphocytic leukemia (ALL). CD7 crosslinking induces a calcium flux in T lymphocytes, presumably as a result of cytoplasmic domain association with PI3-kinase. CD7 co-stimulation can induce cytokine secretion and modulate cellular adhesion. A ligand of CD7, epithelial cell secreted protein K12, is produced in thymus to regulate thymocyte signaling and cytokine release. In lung microvascular endothelial cells CD7 serves as an IgM Fc receptor. Expression of CD7 is an important marker used in leukemia diagnostics.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.




Figure 1: Flow cytometry of PBMC using anti- h CD7 purified antibody