

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1139: Anti-CD71 / Transferrin Receptor Monoclonal Antibody (Clone:MEM-75)

Clonality: Monoclonal Clone Name: MEM-75 Application: **FACS** Reactivity: Human **TFRC** Gene: 7037 Gene ID: **Uniprot ID:** P02786 Format: Purified **TFRC Alternative Name:** Isotype: Mouse IgG1

Immunogen Information: NALM-6 human pre-B cell line

Description

CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathrin-mediated endocytosis. Acidification of endosomes by vesicular membrane proton pumps leads to dissociation of iron ions, whereas transferrin (apotransferrin) remains associated with CD71 and recycles to the cell surface, where it is released upon exposure to normal pH. CD71 is also involved in uptake of non-transferrin bound iron.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow cytometry: Recommended dilution: 0.5-4 µg/ml.




Figure 1: Immunofluorescence staining of CD71 in human HeLa cell line using anti-CD71 (MEM-75; green). Actin cytoskeleton decorated by phalloidin (red) and cell nuclei stained with DAPI (blue).