

### 30-1408: Anti-CD79b Monoclonal Antibody (Clone:HM79)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	HM79
<b>Application :</b>	FACS, IP, WB, ICC
<b>Reactivity :</b>	Mouse
<b>Gene :</b>	Cd79b
<b>Gene ID :</b>	15985
<b>Uniprot ID :</b>	P15530
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Cd79b,Igb
<b>Isotype :</b>	Hamster IgG2
<b>Immunogen Information :</b>	Purified CD79a/b (alpha/beta) dimers from WEHI231 cells

#### Description

CD79b (Ig beta, B29) forms disulfide-linked heterodimer with CD79a (Ig alpha, MB1). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

#### Application Note

**Flow Cytometry** *Recommended dilution:* 1 Åµg/ml

**Immunoprecipitation Western Blotting Immunocytochemistry**