

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1178: Anti-CD80 / B7-1 Monoclonal Antibody (Clone:MEM-233)

Clonality: Monoclonal **Clone Name:** MEM-233 Application: FACS. IP Reactivity: Human Gene: CD80 Gene ID: 941 **Uniprot ID:** P33681 Format: **Purified** 

Alternative Name: CD80,CD28LG,CD28LG1,LAB7

**Isotype:** Mouse IgG1

Immunogen Information: Extracellular domain of human CD80 fused to human IgG1(Fc)

## **Description**

CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.

## **Product Info**

Amount: 0.1 mg

**Purification:** Purified by protein-A affinity chromatography

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Flow Cytometry Recommended dilution:1-10 Ã[βμg/ml Immunoprecipitation