

## 30-1923: Anti-CD89 Monoclonal Antibody (Clone:A59)-FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	A59
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human, Non-Human Primates
<b>Conjugate :</b>	FITC
<b>Gene :</b>	FCAR
<b>Gene ID :</b>	2204
<b>Uniprot ID :</b>	P24071
<b>Alternative Name :</b>	FCAR, CD89, IgA Fc receptor
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Ag8.653 myeloma cells

### Description

CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89 is expressed on granulocytes, monocytes, macrophages, dendritic cells and myeloid cell lines. Its expression is upregulated in presence of IgA immune complexes, stimulators (such as LPS, PMA), TNF alpha, IL1 beta or GM-CSF, and it is downregulated in presence of TGF beta and suramin. Binding of IgA-opsonized targets to CD89 leads to phagocytic and cytotoxic processes of the immunologic defense.

### Product Info

<b>Amount :</b>	100 tests
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of CD89 in human peripheral blood with anti-CD89 (A59) FITC.