

## 30-1623: Anti-CD8a Monoclonal Antibody (Clone:53-6.7)-Low Endotoxin

Clonality :	Monoclonal
Clone Name :	53-6.7
Application :	FACS
Reactivity :	Mouse
Gene :	Cd8a
Gene ID :	12525
Uniprot ID :	P01731
Format :	Low Endotoxin
Alternative Name :	Cd8a,Lyt-2,Lyt2,Ly-2, Ly-B, Ly-35, Lyt-2, BB154331
Isotype :	Rat IgG2a
Immunogen Information : Mouse spleen cells	

## Description

The CD8a (CD8 alpha) subunit of CD8 T cell coreceptor is expressed in CD8 alpha/beta heterodimers on majority of MHC Irestricted conventional T cells and thymocytes and in CD8 alpha/alpha homodimers on subsets of memory T cells, intraepithelial lymphocytes, NK cells, macrophages and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 alpha/beta but not alpha/alpha dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

## **Product Info**

Amount :	0.1 mg
Purification :	Purified by protein-G affinity chromatography
Storage condition :	Store at 2-8°C. Do not freeze.

## **Application Note**

Flow Cytometry Recommended dilution:1.5 µg/ml

**Immunoprecipitation Immunohistochemistry Immunohistochemistry (frozen sections)** *Recommended dilution:* 1:1000

Application note: Formaldehyde fixation is not recommended, acetone fixation is preferred.

**Immunocytochemistry Functional Application** Isolation and depletion of CD8+ cells, blocking of cytotoxicity, inhibition of CD8+ T cell proliferation.