

30-1373: Anti-Cyclin D1 Monoclonal Antibody (Clone:CD1.1)

Clonality :	Monoclonal
Clone Name :	CD1.1
Application :	FACS
Reactivity :	Human
Gene :	CCND1
Gene ID :	595
Uniprot ID :	P24385
Format :	Purified
Alternative Name :	CCND1, DIP1, GCIP, HHM
Isotype :	Mouse IgG1
Immunogen Information :	Purified cyclin D1 protein

Description

Cyclin D1 (PRAD1, Bcl-1) is a cytoplasmic and nuclear protein, which is synthesized during G1 phase and assembles with either cyclin-dependent kinase 4 (CDK4) or CDK6 in response to growth factor stimulation. D-type cyclin-CDK complexes act to inactivate the growth-suppressive function of the Rb protein through its phosphorylation, and titrate CDK inhibitors such as p21Cip1 and p27Kip1. Whereas during G1 phase cyclin D1 accumulates in the nucleus, it translocates into the cytoplasm during S phase. Without growth factor-mediated stimulation cyclin D1 is unstable, and undergoes ubiquitin-mediated degradation, which is triggered by its phosphorylation. Cyclin D1 destabilization participates in G1/S phase arrest.

Product Info

Amount :	0.1 mg
Purification :	Purified by protein-A affinity chromatography
Content :	Formulation: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry *Application note:* Membrane permeabilization is required.

Immunoprecipitation *Recommended dilution:* 1 µg/ml

Western Blotting *Recommended dilution:* 1 µg/ml

Immunohistochemistry *Pretreatment:* Heat treatment, sodium citrate buffer pH 6.0.

Immunohistochemistry (frozen sections) *Recommended dilution:* 2 µg/ml

Positive tissue: colon

Immunocytochemistry *Recommended dilution:* 1 µg/ml

ELISA




Figure 1: Separation of MCF-7 cells stained using anti-Cyclin D1 (CD1.1) purified antibody (GAM APC, red-filled) from MCF-7 cells stained using mouse IgG1 isotype control (MOPC-21) purified antibody (GAM APC, black-dashed) in flow cytometry analysis (intracellular staining).