

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1829: Biotin Conjugated, Anti-Cytokeratin 18 Monoclonal Antibody (Clone:DA-7)

Clonality: Monoclonal

Clone Name: DA-7

**Application:** IP, WB, IHC, ICC, ELISA

Reactivity: Human
Conjugate: Biotin
Gene: KRT18
Gene ID: 3875
Uniprot ID: P05783

**Alternative Name:** KRT18,CYK18,PIG46

**Isotype:** Mouse IgG1

Immunogen Information: Human breast carcinoma cell line PMC-42.

## **Description**

Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families.Cytokeratins 18 belongs to type I family (acidic cytokeratins).

## **Product Info**

Amount: 0.1 mg

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Western blotting: Recommended dilution: 1-2 Ã\|\hat{A}\|\hat{4}g/ml.




Figure-1: Western blotting analysis of human cytokeratin 18 using mouse monoclonal antibody DA-7 on lysates of HT-29 cell line and MOLT-4 cell line (cytokeratin non-expressing cell line; negative control) under non-reducing and reducing conditions. Nitrocellulose membrane was probed with 2 µg/ml of biotinylated mouse anticytokeratin 18 monoclonal antibody followed by IRDye800-conjugated streptavidin. Cytokeratin 18 was detected at approximately 46 kDa.