

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1920: FITC Conjugated Anti-p53 Monoclonal Antibody (Clone:BP53-12)

Clonality: Monoclonal **Clone Name:** BP53-12 **FACS** Application: Reactivity: Human Conjugate: FITC Gene: TP53 Gene ID: 7157 **Uniprot ID:** P04637 **Alternative Name:** TP53,P53 Isotype: Mouse IgG2a

Immunogen Information: Bacterially expressed full-length wild-type p53

Description

The tumour suppressor protein p53 is a key element of intracellular anticancer protection. It mediates cell cycle arrest or apoptosis in response to DNA damage or to starvation for pyrimidine nukleotides. It is up-regulated in response to these stress signals and stimulated to activate transcription of specific genes, resulting in expression of p21waf1 and other proteins involved in G1 or G2/M arrest, or proteins that trigger apoptosis, such as Bcl-2. The structure of p53 comprises N-terminal transactivation domain, central DNA-binding domain, oligomerisation domain, and C-terminal regulatory domain. There are various phosphorylation sites on p53, of which the phosphorylation at Ser15 is important for p53 activation and stabilization.

Product Info

Amount: 0.1 mg

Storage condition: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Confocal microscopy of human HeLa cells using anti-p53 (BP53-12; FITC). The expression of p53 protein was enhanced by intercalating reagent. Cells were fixed and permeabilized before incubation with the p53-FITC MAb.Photo provided by Dr. Hodny, Inst. of Experimental Medicine, Prague, Czech Republic



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Intracellular staining of p53 in RAMOS cells with anti-p53 (BP53-12) FITC.