

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1425: Anti-TCR alpha/beta Monoclonal Antibody (Clone:IP26)

Clonality: Monoclonal

Clone Name: IP26
Application: FACS
Reactivity: Human

Gene : Pre T-Cell Receptor alpha beta

Gene ID: 171558 Uniprot ID: Q6ISU1 Format: Purified

Alternative Name: Pre T-Cell Receptor alpha beta

Isotype: Mouse IgG1

Description

The antigen-specific T cell receptor (TCR) is composed of either alpha and beta subunit, or gamma and delta subunit. Majority of T cells present in the blood, lymph and secondary lymphoid organs express TCR alpha/beta heterodimers, whereas the T cells expressing TCR gamma/delta heterodimers are localized mainly in epithelial tissues and at the sites of infection. The subunits of TCR heterodimers are covalently bonded and in the endoplasmic reticulum they associate with CD3 subunits to form functional TCR-CD3 complex. Lack of expression of any of the chains is sufficient to stop cell surface expression.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution:2-4 µg/ml Positive control:human peripheral blood T cells




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR alpha/beta (IP26) purified antibody



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of human TCR alpha/beta positive lymphocytes (red-filled) from neutrofil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of peripheral whole blood stained using anti-human TCR alpha/beta (IP26) purified antibody