

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2002: Mouse Monoclonal Antibody To N-Cadherin (Clone: CDH2/1573)

Clonality: Monoclonal Clone Name: CDH2/1573

Application: IHC

Reactivity: Mouse, Human

 Gene :
 CDH2

 Gene ID :
 1000

 Uniprot ID :
 P19022

**Isotype:** Mouse IgG1, kappa

Immunogen Information: Recombinant human CDH2 intracellular domain was used as thye immunogen

## **Description**

Recognizes a protein of ~140kDa, identified as N-Cadherin (NCAD), also known as CD325. N-cadherin is a 140 kDa protein belonging to a family of transmembrane molecules that mediate calcium-dependent intercellular adhesion. Cadherins are involved in controlling morphogenetic movements during development and regulate cell surface adhesion through homotypic adhesion with the same cadherin species. Expression of N-cadherin has been reported on a variety of normal tissues including neuronal, endothelial and muscle cells, and a subpopulation of early hematopoietic progenitor cells. Results aid in the classification of malignant non-carcinomatous neoplasms including mesotheliomas, chordomas, synovial sarcomas, malignant melanomas, epithelioid sarcomas, epithelioid angiosarcomas, clear cell sarcomas as well as serous and endometrioid tumors of the ovary have been demonstrated to be N-cadherin positive, whereas mucinous tumors are negative. Other N-cadherin-positive neoplasms include renal cell carcinomas and some variant breast tumors, including medullary breast carcinomas and sarcomatoid metaplastic breast carcinomas.

## **Product Info**

**Amount :** 100 μg

**Purification :** Affinity Chromatography

Content: 200 µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10 mM PBS with

0.05% BSA & 0.05% azide.

Storage condition:

Storage condition:

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.


## **Application Note**

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes)



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Immunochemical analysis of FFPE mouse heart with Cadherin 2 antibody (clone CDH2/1573).