

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-4199: Monoclonal Antibody to mouse MHC Class II (I-A/I-E) (Clone: M5/114)-NALE

Clonality: Monoclonal Clone Name: M5/114

Application: Functional Assay,IHC,FACS,WB,IF

Reactivity: Mouse

Format: Low Endotoxin, Azide Free

Isotype: Rat IgG2b, k

Immunogen Information: Activated C57BL/6 mouse spleen cells was used as Immunogen

Description

MHC (major histocompatibility complex) class II molecules are transmembrane glycoproteins expressed on the surface of professional antigen-presenting cells, such as macrophages, dendritic cells and B cells. Before their exposition on the cell surface, the MHC class II molecules react with endocytosed exogenous antigens, which are then presented to the T cells. The antigen-binding grove between MHC class II alpha and beta chain is open at both ends and is 15-24 amino acid residues long.

Product Info

Amount : $25 \mu g / 100 \mu g$ Purification : Protein G

Content : PBS, pH 7.0 Contains no stabilizers or preservatives

Storage condition : The antibody solution should be stored undiluted at 4°C, and protected from prolonged

exposure to light. Do not freeze.

Application Note

<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay Functional assays, Immunofluorescence, Western blot, Immunoprecipitation, Flow cytometry




Figure-1: Cell surface flow analysis of mouse MHC Class II on C57 mice splenocytes using 0.5 μ g/10^6 cells of MHC class II antibody (Clone: M5/114). Green represents isotype control; red represents anti-mouse MHC class II antibody. mouse anti-rat Fitc conjugate was used as secondary antibody.