

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 42-1340: Anti-HSP40, YDJ1 Monoclonal Antibody (Clone: 1G10.H8) - PerCP

Clone Name : Monoclonal
Clone Name : 1G10.H8
Application : WB,IP,ELISA

 Reactivity:
 Yeast

 Conjugate:
 PerCP

 Gene:
 YDJ1

 Gene ID:
 855661

 Uniprot ID:
 P25491

Alternative Name: YDJ1,MAS5,YNL064C,N2418,YNL2418C

**Isotype:** Mouse IgG1 Kappa

Immunogen Information: Full length protein yeast HSP40 (YDJ1)

## **Description**

Human HSP40/DnaJ proteins comprise a large protein family, members of which feature the J domain (named after the bacterial DnaJ protein). The J-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain. There are two main types of HSP40; type 1 DNAJ proteins including HDJ2 and yeast YdjI; type II includes yeast Sis1 and human Hdj1. Whereas type I possesses a zinc finger domain which helps in the function of protein folding, type II does not. Members of the HSP40/DnaJ family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins. The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1.

## **Product Info**

**Amount :** 100 μg

Purification: Protein G Purified

**Content:** 50% glycerol, 0.09% sodium azide

Storage condition: Store the antibody at 4°C

## **Application Note**

WB (1:2000); optimal dilutions for assays should be determined by the user.




Figure1: Mouse Anti-Hsp40 Antibody [1G10.H8] used in Western Blot (WB) on Yeast Cell Ivsates