

## 32-20007: Recombinant Human ApoE2(Discontinued)

**Alternative Name :** Apolipoprotein E2

### Description

**Source: E.coli**

ApoE belongs to a group of proteins that bind reversibly with lipoprotein and play an important role in lipid metabolism. In addition to facilitating solubilization of lipids, these proteins help to maintain the structural integrity of lipoproteins, serve as ligands for lipoprotein receptors, and regulate the activity of enzymes involved in lipid metabolism. Significant quantities of ApoE are produced in the liver and brain, and to some extent in almost every organ. ApoE is an important constituent of all plasma lipoproteins. Its interaction with specific ApoE receptor enables uptake of chylomicron remnants by liver cells, which is an essential step during normal lipid metabolism. It also binds with the LDL receptor (apo B/E). Defects in ApoE are a cause of hyperlipoproteinemia type III. ApoE exists in three major isoforms; E2, E3, and E4, which differ from one another by a single amino-acid substitution. Compared with E3 and E4, E2 exhibits the lowest receptor binding affinity. E2 allele carriers had significantly lower levels of total cholesterol, low-density lipoprotein cholesterol, and non-high-density lipoprotein cholesterol, as well as increased ApoE levels. Recombinant Human ApoE2 is a 34.3 kDa protein containing 300 amino acid residues.

### Product Info

**Amount :** 100 µg / 500 µg

**Purification :** Purity:  $\geq 90\%$  by SDS-PAGE gel and HPLC analyses.

**Amino Acid :** MKVEQAVETE PEPELRQQTE WQSGQRWELA LGRFWDYLRW VQTLSEQVQE ELLSSQVTQE  
LRALMDETMK ELKAYKSELE EQLTPVAEET RARLSKELQA AQARLGADME DVCGRVLQYR  
GEVQAMLGQS TEELRVRLAS HLRKLRKRL RDADDLQKCL AVYQAGAREG AERGLSAIRE RLGPLVEQGR  
VRAATVGSLA GQPLQERAQA WGERLRARME EMGSRTDRDL DEVKEQVAEV RAKLEEQAQQ  
IRLQAEAFQA RLKSWFEPLV EDMQRQWAGL VEKVQAAVGT SAAPVPSDNH