# **w** abeomics

# 32-20085: Recombinant Human sFas Receptor(Discontinued)

Alternative Name : soluble Fas receptor (sFasR), TNFRSF6, CD95, Apo I, Fas Antigen

## Description

**Source:E.coli**Fas and Fas Ligand (FasL) belong to the TNF superfamily, and are type I and type II transmembrane proteins, respectively. Binding of FasL to Fas triggers apoptosis in Fas-bearing cells. The mechanism of apoptosis involves recruitment of pro-caspase 8 through an adaptor molecule called FADD, followed by processing of the pro-enzyme into active forms. These active caspases then cleave various cellular substrates, leading to the eventual cell death. sFasR is capable of inhibiting FasL-induced apoptosis by acting as a decoy receptor that serves as a sink for FasL. The full length Fas (receptor) is a 319 amino acid type I transmembrane protein, which contains a 157 amino acid extracellular domain, a 17 amino acid transmembrane domain, and a 145 amino acid cytoplasmic domain. Recombinant Human soluble Fas (sFas Receptor) is a 157 amino acid polypeptide (17.6 kDa) corresponding to the TNFR-homologous cysteine-rich extracellular Fas domain.

### **Product Info**

 

 Amount :
 5 μg / 20 μg

 Purification :
 Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

 Content :
 This recombinant protein is supplied in lyophilized form.

 Amino Acid :
 MRLSSKSVNA QVTDINSKGL ELRKTVTTVE TQNLEGLHHD GQFCHKPCPP GERKARDCTV NGDEPDCVPC QEGKEYTDKA HFSSKCRRCR LCDEGHGLEV EINCTRTQNT KCRCKPNFFC NSTVCEHCDP CTKCEHGIIK ECTLTSNTKC KEEGSRS

#### **Application Note**

The  $\tilde{A} \equiv \tilde{A} \equiv D_{50} \tilde{A} \equiv \tilde{A}$  was determined by its ability to inhibit the cytotoxicity of Jurkat cells is between 10-15  $\tilde{A} \equiv \tilde{A} \mu g/ml$  in the presence of 2ng/ml of hFasL.