

## 32-20094: Recombinant Murine FGF-basic(Discontinued)

**Reactivity :** Human , mouse, Rabbit, rat

**Alternative Name :** Fibroblast Growth Factor-basic, FGF-2, HBGF-2, Prostatropin

### Description

**Source:** **E.coli** FGF-basic is one of 23 known members of the FGF family. Proteins of this family play a central role during prenatal development, postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF-basic is a non-glycosylated, heparin-binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland, liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4. Recombinant Murine FGF-basic is a 16.3 kDa protein consisting of 145 amino acid residues.

### Product Info

**Amount :** 10 µg / 50 µg

**Purification :** Purity:  $\geq 95\%$  by SDS-PAGE gel and HPLC analyses.

**Content :** This recombinant protein is supplied in lyophilized form.

**Amino Acid :** PALPEDGGAA FPPGHFKDPK RLYCKNGGFF LRIHPDGRVD GVREKSDPHV KLQLQAEERG VVSIKGVCAN  
RYLAMKEDGR LLASKCVTEE CFFFERLESN NYNTYRSRKY SSWYVALKRT GQYKLGSKTG PGQKAILFLP  
MSAKS

### Application Note

Determined by a cell proliferation assay using Balb/c 3T3 cells. The expected  $ED_{50}$  is  $\leq 1.0$  ng/ml, corresponding to a specific activity of  $\geq 1 \times 10^6$  units/mg.