

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-20176: Recombinant Murine IFN-Gamma(Discontinued)

Reactivity: Cow, Human, Rabbit, Rat

Alternative Name: Immune Interferon, type II interferon, T cell interferon, MAF

Description

Source:E.coliIFN-Gamma is an acid-labile interferon produced by CD4 and CD8 T lymphocytes, as well as activated NK cells. IFN-Gamma receptors are present in most immune cells, which respond to IFN-Gamma signaling by increasing the surface expression of class I MHC proteins. This promotes the presentation of antigen to T-helper (CD4+) cells. IFN-Gamma signaling in antigen-presenting, cells and antigen-recognizing B and T lymphocytes, regulates the antigen-specific phases of the immune response. Additionally, IFN-Gamma stimulates a number of lymphoid cell functions, including the anti-microbial and anti-tumor responses of macrophages, NK cells, and neutrophils. Human IFN-Gamma is species-specific and is biologically active only in human and primate cells. Recombinant Murine IFN-Gamma is a 15.6 kDa protein containing 134 amino acid residues.

Product Info

Amount : 20 μg / 100 μg

Purification: Purity: >= 98% by SDS-PAGE gel and HPLC analyses. **Content**: This recombinant protein is supplied in lyophilized form.

Amino Acid: MHGTVIESLE SLNNYFNSSG IDVEEKSLFL DIWRNWQKDG DMKILQSQII SFYLRLFEVL

KDNQAISNNI SVIESHLITT FFSNSKAKKD AFMSIAKFEV NNPQVQRQAF NELIRVVHQL

LPESSLRKRK RSRC

Application Note

Determined by its ability to inhibit the proliferation of murine WEHI-279 cells. The expected ED_{50} is <= 0.2 ng/ml, corresponding to a specific activity of >= 5 x 10^6 units/mg.