9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-20177: Animal-Free Recombinant Murine IFN-Gamma(Discontinued)

Reactivity : Human, Mouse
Alternative Name : Immune Interferon, type II interferon, T cell interferon, MAF

Description

Source:E.coliIFN-Gamma is an acid-labile interferon produced by CD4 and CD8 T lymphocytes, as well as activated NK cells.Â IFN-Gamma receptors are present in most immune cells, which respond to IFN-Gamma signaling by increasing the surface expression of class I MHC proteins.Â This promotes the presentation of antigen to T-helper (CD4+) cells.Â IFN-Gamma signaling in antigen-presenting, cells and antigen-recognizing B and T lymphocytes, regulates the antigen-specific phases of the immune response.Â Additionally, IFN-Gamma stimulates a number of lymphoid cell functions, including the anti-microbial and anti-tumor responses of macrophages, NK cells, and neutrophils.Â Human IFN-Gamma is species-specific and is biologically active only in human and primate cells.Â Recombinant Murine IFN-Gamma is a 15.6 kDa protein containing 134 amino acid residues.

Product Info

Amount : 20 µg / 100 µg
Purification : Purity: >= 98% by SDS-PAGE gel and HPLC analyses.
Content : This recombinant protein is supplied in lyophilized form.
Amino Acid : MHGTVIESLE SLNNYFNSSG IDVEEKSLFL DIWRNWQKDG DMKILQSQII SFYLRLFEVL KDNQAISNNI SVIESHLITT FFSNSKAKKD AFMSIAKFEV NNPQVQRQAF NELIRVVHQL LPESSLRKRK RSRC

Application Note

Determined by its ability to inhibit the proliferation of murine WEHI-279 cells. The expectedÃ□Â ED_{50}Ã□Â is <= 0.2 ng/ml, corresponding to a specific activity of >= 5 x 10^6units/mg.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.