

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-20178: Recombinant Rat IFN-Gamma(Discontinued)

Reactivity: Human, Mouse, Rat

Alternative Name: Immune Interferon, type II interferon, T cell interferon, MAF

## **Description**

**Source:E.coli**IFN-Gamma is an acid-labile interferon produced by CD4 and CD8 T lymphocytes, as well as activated NK cells. IFN-Gamma receptors are present in most immune cells, which respond to IFN-Gamma signaling by increasing the surface expression of class I MHC proteins. This promotes the presentation of antigen to T-helper (CD4+) cells. IFN-Gamma signaling in antigen-presenting cells, and antigen-recognizing B and T lymphocytes, regulates the antigen-specific phases of the immune response. Additionally, IFN-Gamma stimulates a number of lymphoid cell functions, including the anti-microbial and anti-tumor responses of macrophages, NK cells, and neutrophils. Human IFN-Gamma is species-specific and is biologically active only in human and primate cells. Recombinant Rat IFN-Gamma is a 15.6 kDa protein containing 135 amino acid residues.

## **Product Info**

**Amount:** 20 μg / 100 μg

**Purification**: Purity: >= 98% by SDS-PAGE gel and HPLC analyses. **Content**: This recombinant protein is supplied in lyophilized form.

Amino Acid: MQGTLIESLE SLKNYFNSSS MDAMEGKSLL LDIWRNWQKD GNTKILESQI ISFYLRLFEV

LKDNQAISNN ISVIESHLIT NFFSNSKAKK DAFMSIAKFE VNNPQIQHKA VNELIRVIHQ

LSPESSLRKR KRSRC

## **Application Note**

The ED<sub>50</sub>Â as determined by a cytopathic affect inhibition assay with murine L929 cells challenged with EMC virus was  $\leq 0.1$  ng/ml, corresponding to a specific activity of  $\geq 1 \times 10^7$  units/mg.