

## 32-20203: Recombinant Murine IL-1Beta (Discontinued)

Reactivity : Human, Mouse, Pig, Rat

Alternative Name : Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF)

## Description

**Source:E.coli**IL-1Beta is a proinflammatory cytokine produced in a variety of cells, including monocytes, tissue macrophages, keratinocytes, and other epithelial cells. Both IL-1Alpha and IL-1Beta bind to the same receptor and have similar, if not identical, biological properties. These cytokines have a broad range of activities including the stimulation of thymocyte proliferation by inducing IL-2 release, B cell maturation and proliferation, mitogenic FGF-like activity, and the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1Beta is a secreted cytokine, IL-1Alpha is predominantly a cell-associated cytokine. Recombinant Murine IL-1Beta is a 17.5 kDa protein containing 153 amino acid residues.

## **Product Info**

 Amount :
 2 μg / 10 μg

 Purification :
 Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

 Content :
 This recombinant protein is supplied in lyophilized form.

 Amino Acid :
 MVPIRQLHYR LRDEQQKSLV LSDPYELKAL HLNGQNINQQ VIFSMSFVQG EPSNDKIPVA LGLKGKNLYL SCVMKDGTPT LQLESVDPKQ YPKKKMEKRF VFNKIEVKSK VEFESAEFPN WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS

## **Application Note**

The  $\tilde{A} \equiv \tilde{A} \equiv D_{50} \tilde{A} \equiv \tilde{A}$  as determined by the dose-dependent stimulation of murine D10S cells is <= 0.002 ng/ml, corresponding to a specific activity of >= 5 x 108units/mg.