9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-20170: Recombinant Murine SCF(Discontinued)

Reactivity : Human, Monkey, Mouse, Rat
Alternative Name : Stem Cell Factor, c-Kit Ligand, Mast Cell Growth Factor (MGF), Steel Factor

Description

Source:E.coli
SCF is a hematopoietic growth factor that exerts its activity by signaling through the c-Kit receptor. SCF and c-Kit are essential for the survival, proliferation and differentiation of hematopoietic cells committed to the melanocyte and germ cell lineages. Human SCF manifests low activity on murine cells, while murine and rat SCF are fully active on human cells. The human SCF gene encodes for a 273 amino acid transmembrane protein, which contains a 25 amino acid N-terminal signal sequence, a 189 amino acid extracellular domain, a 23 amino acid transmembrane domain, and a 36 amino acid cytoplasmic domain. The secreted soluble form of SCF is generated by proteolytic processing of the membrane anchored precursor. Recombinant Murine SCF is an 18.3 kDa polypeptide containing 165 amino acid residues, which corresponds to the sequence of the secreted soluble form of SCF.

Product Info

Amount : 2 µg / 10 µg
Purification : Purity: >= 98% by SDS-PAGE gel and HPLC analyses.
Amino Acid : MKEICGNPVT DNVKDITKLV ANLPNDYMIT LNYVAGMDVL PSHCWLRDMV IQLSLSLTTL LDKFSNISEG LSNYSIIDKL GKIVDDLVLC MEENAPKNIK ESPKRPETRS FTPEEFFSIF NRSIDAFKDF MVASDTSDCV LSSTLGPEKD SRVSVTKPFM LPPVA

Application Note

TheÃ Â ED_{50} Ã as determined by the dose-dependent stimulation of the proliferation of the human TF-1 cells isÃ <= 10 ng/ml, corresponding to a specific activity ofÃ >= 1×10^{6}

For Research Use Only. Not for use in diagnostic/therapeutics procedures.