

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-20225: Recombinant Rat SDF-1Beta (CXCL12)(Discontinued)

Reactivity: Human, Rat

Alternative Name: Stromal-Cell Derived Factor-1, CXCL12, PBSF

Description

Source: E.coli

SDF-1Alpha and Beta are stromal-derived, CXC chemokines that signal through the CXCR4 receptor. SDF-1Alpha and Beta chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV-suppressive activity in cells expressing the CXCR4 receptor. Human and murine SDF-1 proteins act across species. SDF-1Alpha and Beta contain the four highly conserved cysteine residues present in CXC chemokines. The mature SDF-1Beta protein, produced by an N-terminal truncation of two additional amino acids, after removal of the signal sequence, contains 72 amino acid residues. Recombinant Rat SDF-1Beta is an 8.4 kDa protein containing 72 amino acid residues.

Product Info

Amount: 2 μg / 10 μg

Purification : Purity: >= 95% by SDS-PAGE gel and HPLC analyses.

Amino Acid: KPVSLSYRCP CRFFESHVAR ANVKHLKILN TPNCALQIVA RLKSNNRQVC IDPKLKWIQE

YLDKALNKRL KM

Application Note

Determined by its ability to chemoattract human monocytes using a concentration range of 100.0-200.0 ng/ml.