

32-20229: Animal-Free Recombinant Human TGF-Beta3(Discontinued)

Reactivity : Human, Mouse, Rat

Alternative Name : Transforming Growth Factor-Beta3

Description

Source:E.coli

The three mammalian isoforms of TGF-Beta, TGF-Beta1, Beta2, and Beta3, signal through the same receptor and elicit similar biological responses. They are multifunctional cytokines that regulate cell proliferation, growth, differentiation and motility, as well as synthesis and deposition of the extracellular matrix. They are involved in various physiological processes, including embryogenesis, tissue remodeling and wound healing. They are secreted predominantly as latent complexes, which are stored at the cell surface and in the extracellular matrix. The release of the biologically active TGF-Beta isoform from a latent complex involves proteolytic processing of the complex and/or induction of conformational changes by proteins such as thrombospondin-1. The physiological role of TGF-Beta3 is still unknown, but its expression pattern suggests a role in the regulation of certain development processes. Recombinant Human TGF-Beta3 is a 25.0 kDa protein composed of two identical 112 amino acid polypeptide chains linked by a single disulfide bond.

Product Info

Amount : 2 µg / 10 µg

Purification : Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

Amino Acid : ALDTNYCFRN LEENCCVRPL YIDFRQDLGW KVVHEPKGY ANFCSGPCPY LRSADTTHT
VLGLYNTLNP EASASPCCVP QDLEPLTILY YVGRTPKVEQ LSNMVKSCCK CS

Application Note

Determined by TGF-Beta3's ability to inhibit the mouse IL-4-dependent proliferation of mouse HT-2 cells. The expected ED_{50} is ≈ 0.05 ng/ml, corresponding to a specific activity of 2×10^6 IU/mg.