

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

# 32-20382: Animal-Free Recombinant Human IL-22(Discontinued)

Alternative Name: Interleukin-22, IL-TIF, Cytokine ZCYTO18

# **Description**

#### Source: E.coli

IL-22 is a member of the IL-10 family of regulatory cytokines, which includes IL-10, IL-19, IL-20, IL-22, IL-24 and IL-26. Members of this family share partial homology in their amino acid sequences, but they are dissimilar in their biological functions. Produced by T lymphocytes, IL-22 inhibits IL-4 production by Th2 cells, and induces acute phase reactants in the liver and pancreas. IL-22 signals through a receptor system consisting of IL-10RBeta /CRF2-4 and IL-22R, both of which are members of the class II cytokine-receptor family. Recombinant Human IL-22 is a 33.6 kDa, non-disulfide-linked, homodimeric protein consisting of two 147 amino acid polypeptide chains.

### **Product Info**

Amount:  $2 \mu g / 10 \mu g$ 

Purification: Purity:>= 98% by SDS-PAGE gel and HPLC analyses.Content: This recombinant protein is supplied in lyophilized form.

Amino Acid: MAPISSHCRL DKSNFQQPYI TNRTFMLAKE ASLADNNTDV RLIGEKLFHG VSMSERCYLM

KQVLNFTLEE VLFPQSDRFQ PYMQEVVPFL ARLSNRLSTC HIEGDDLHIQ RNVQKLKDTV

KKLGESGEIK AIGELDLLFM SLRNACI

# **Application Note**

Measured by its ability to induce IL-10 secretion in COLO 205 (human colon carcinoma cells). The expected  $ED_{50}$  for this effect is 0.7- 1.0 ng/ml.