w abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

32-20396: Animal-Free Recombinant Human M-CSF(Discontinued)

Reactivity : Human, Monkey, Mouse, Rat

Alternative Name : Macrophage Colony Stimulating Factor, CSF-1, MGI-IM

Description

Source:E.coliM-CSF is a potent hematopoietic factor produced by a variety of cells, including lymphocytes, monocytes, fibroblasts, endothelial cells, myoblasts and osteoblasts. It is a key regulator of cellular proliferation, differentiation, and survival for blood monocytes, tissue macrophages, and their respective progenitor cells. M-CSF has been shown to play important roles in modulating dermal thickness and fertility. M-CSF is clinically used in the treatment of infection, malignancies and atherosclerosis. It facilitates hematopoietic recovery after bone marrow transplantation. Human M-CSF is reactive in murine systems, but the murine molecule exhibits no activity on human cells. Recombinant Human M-CSF is a 36.8 kDa homodimeric protein consisting of two 159 amino acid polypeptide subunits.

Product Info

 Amount :
 2 μg / 10 μg

 Purification :
 Purity:>= 98% by SDS-PAGE gel and HPLC analyses.

 Content :
 This recombinant protein is supplied in lyophilized form.

 Amino Acid :
 MEEVSEYCSH MIGSGHLQSL QRLIDSQMET SCQITFEFVD QEQLKDPVCY LKKAFLLVQD IMEDTMRFRD NTPNAIAIVQ LQELSLRLKS CFTKDYEEHD KACVRTFYET PLQLLEKVKN VFNETKNLLD KDWNIFSKNC NNSFAECSSQ GHERQSEGS

Application Note

The $\tilde{A} \equiv \tilde{A} \equiv \tilde{A}_{50} \tilde{A} \equiv \tilde{A}$ was determined by the dose-dependent stimulation of the proliferation of murine M-NFS-60 cells is <=1.0 ng/ml, corresponding to a specific activity of>= 1 x 10⁶ $\tilde{A} \equiv \tilde{A}$ units/mg. $\tilde{A} \equiv \tilde{A}$